

## Guideline-adherent antithrombotic treatment improves outcomes in patients with atrial fibrillation:

Mazurek, Michal; Shantsila, Eduard; Lane, Deirdre; Wolff, Andreas; Proietti, Marco; Lip, Gregory

DOI:

[10.1016/j.mayocp.2017.05.023](https://doi.org/10.1016/j.mayocp.2017.05.023)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Mazurek, M, Shantsila, E, Lane, D, Wolff, A, Proietti, M & Lip, G 2017, 'Guideline-adherent antithrombotic treatment improves outcomes in patients with atrial fibrillation: Insights from the community-based Darlington atrial fibrillation registry', *Mayo Clinic Proceedings*, vol. 92, no. 8, pp. 1203-1213.  
<https://doi.org/10.1016/j.mayocp.2017.05.023>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.


### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.


If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Supplemental Figure 1 One-Year Stroke Rates in Relation to Antithrombotic Guideline Adherence

**Panel A** Entire population (unselected study cohort)



**Panel B** High-risk cohort (CHA<sub>2</sub>DS<sub>2</sub>-VASc ≥2)



## Figure Legend

### **Supplemental Figure 1** One-Year Stroke Rates in Relation to Antithrombotic Guideline Adherence

**Panel A** Entire population (unselected study cohort)

**Panel B** High-risk cohort (CHA<sub>2</sub>DS<sub>2</sub>-VASc  $\geq 2$ )

CHA<sub>2</sub>DS<sub>2</sub>-VASc = congestive heart failure, hypertension, age  $\geq 75$  years, diabetes mellitus, stroke/transient ischemic attack (TIA), vascular disease, age 65-74 years, sex category (female); OAC = oral anticoagulant.

Some percentages may not sum up to total due to rounding.